

**Description of Map Units**

QUATERNARY SYSTEM  
 HOLOCENE

- Hua** **Holocene undifferentiated alluvium**—undifferentiated deposits of small upland streams; alluvial deposits of minor streams and creeks, of varying textures, filling valleys incised into older deposits.
- Hb** **Backswamp deposits**—fine-grained Holocene deposits of rivers, underlying the flood basins between meander belts.
- Hod** **Ouachita River distributary deposits**—sandy and silty sediments occupying abandoned courses of a relict distributary system of the Ouachita River. In the Monroe area the sediments comprise yellowish to orangish brown silty very fine sand with varying though relatively small proportions of admixed clay.
- Hol** **Ouachita River natural levee deposits**—deposits forming low natural levees flanking the meander belts of the Ouachita River. Where observed in the Monroe area the sediments comprise grayish brown silty clay with well developed soil structure.
- Hocs** **Ouachita River crevasse splay deposits**—sandy and silty sediments forming fanlike crevasse splays that originate from the Ouachita River. Where observed in the Monroe area the sediments comprise interlaminated gray-brown silt and organic-rich, dark clayey silt.
- Hom** **Ouachita River meander-belt deposits**—point bar deposits underlying meander belts of the Ouachita River.
- Hcr** **River channel remnants**—sinuous tonal patterns interpreted to be abandoned river channels, buried beneath backswamp and natural levee deposits.

PLEISTOCENE

- PLAINE ALLOGROUP**
- Plu** **Upper Lapine alloformation**—stratigraphically higher sequence underlying the lower of two depositional surfaces of the Lapine alloformation. Silty to sandy clay, and clayey to silty very fine to fine sand, of grayish to dark gray coloration with orange-brown to red mottles. May be veneered locally by silty colluvium.
- Pir** **Lapine alloformation, relict beach ridge**—relict shoreline ridges formed on depositional surfaces of the Lower Lapine alloformation. Grayish very fine to medium sand, in places clayey and/or gravelly, weathering to tan, yellowish brown, orangish brown, or reddish brown coloration. The texture typically becomes clayier away from the ridge crests. The upper 1 to 2 m may consist of a reddish brown weathering zone in places.
- Plm** **Lapine alloformation meander-belt deposits**—sandy point bar deposits underlying the abandoned course of a meander belt formed on the depositional surface of the upper Lapine alloformation. Grayish clayey very fine to fine sand with orange-brown mottles.
- Pli** **Lower Lapine alloformation**—stratigraphically lower sequence underlying the higher of two depositional surfaces of the Lapine alloformation. Tan to light gray fine to very fine sand, in places clayey and/or gravelly, weathering to yellowish brown or orangish brown coloration. Ironstone nodules are common and range in size up to 0.6 m. Contains some admixed dark brown organic material in places. May be veneered locally by silty colluvium.

**Open Water, Inundated Area, Wetland**

- Contact**—includes inferred contacts.
- Railroads**
- Streams**
- Topographic Contours**

References:

Andersen, H. V., 1960, Geology of Sabine Parish: Louisiana Department of Conservation, Louisiana Geological Survey, Geological bulletin no. 34, 164 p. plus plates (includes one 1:62,500-scale geologic map).

Andersen, H. V., 1993, Geology of Natchitoches Parish: Louisiana Geological Survey, Geological bulletin no. 44, 227 p. plus plates (includes one 1:62,500-scale geologic map).

Produced and published by the Louisiana Geological Survey  
 3079 Energy, Coast & Environment Building, Louisiana State University  
 Baton Rouge, LA 70803 • 225/578-5320 • www.lgs.lsu.edu

This geologic map was funded in part by the USGS National Cooperative  
 Geologic Mapping Program under STATEMAP award number 04HQAG0073



Copyright ©2006 and 2021 by the Louisiana Geological Survey

Geology by: Richard P. McCulloh and Paul V. Heinrich


GIS Compilers: R. Hampton Peele, Jayanth Ramachandran, and Jomish George

Revision GIS: Robert Pausell

Cartography by: Robert Pausell



1	2	3	1 West Monroe North
4	5	2 Monroe North	2 Monroe North
6	7	3 West Monroe South	3 West Monroe South
8	8	4 West Monroe South	4 West Monroe South
		5 Crew Lake	5 Crew Lake
		6 Luma	6 Luma
		7 Boico	7 Boico
		8 Gilleville	8 Gilleville



MONROE SOUTH, LA  
 2018

This research is supported by the U. S. Geological Survey, National Cooperative Geologic Mapping Program, under USGS award number G20AC00239, 2020. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U. S. Government or the state of Louisiana. This map was produced to conform with the National Geospatial Program US Topo Product Standard, 2011.

This map has been carefully prepared from the best existing sources available at the time of preparation. However, the Louisiana Geological Survey and Louisiana State University do not assume responsibility or liability for any reliance thereon. This information is provided with the understanding that it is not guaranteed to be correct or complete, and conclusions drawn from such data are the sole responsibility of the user. These geologic quadrangles are intended for use at the scale of 1:24,000. A detailed on-the-ground survey and analysis of a specific site may differ from these maps.